

DAVID L. LIMAURO, MD MARK A. CEDAR, DO NICHOLAS BELLICINI, DO XUONG LU, MD JUNGMIN LEE, MD LISA A. OLIVA, DO THERESA SCHUERLE, DO BRIDGER CLARKE, MD Pittsburgh Gastroenterology Associates 2589 Boyce Plaza Road Pittsburgh, PA 15241 PHONE: 412.232.8104 FAX: 412.281.1898 www.pghgastro.com

Name:		
You are scheduled for a consultation a	and <i>Upper End</i>	oscopy or EGD with possible biopsy
and/or polypectomy on	at	Please arrive at the
hospital or surgery center no later the	en:	Your procedure is scheduled at:
South Hills Endoscopy Center. the building. Parking is free.	Report to the recep	otion desk in the main lobby in the front of
Mercy Hospital. Report to the End floor of Mercy Tower. Reduced park	1.	al Procedures Unit located on the second he hospital lot.
	rea around the hos	Family Center located on the sixth floor. A pital. Call transportation services through nents.
<u>Spartan Surgicenter</u> . Report to to Surgicenter Building. Parking is free	-	ocated on the ground floor of the Spartan
Penn Highlands Mon Valley . Re	eport to same day s	urgery on first floor.
Waterfront Surgery Center. Rep	port to the reception	n desk in the main lobby.
McCandless Endoscopy Center. the building.	Report to the rece	eption desk in the main lobby in the back of

<u>Please call 48 hours in advance</u> <u>if you are unable to keep your appointment</u>

- Since you will be sedated for this exam, **you CANNOT drive** or operate machinery for 12-24 hours following the test. **You must be accompanied by a companion and make arrangements for transportation** or you will not be given the sedation.
- It is very important that you strictly follow the instruction:

• As Soon As You Get This Instruction:

- o <u>Check your medication</u> and make necessary adjustments if you take blood thinners or diabetic drugs (see below).
- o Make arrangements for transportation since you are not allowed to drive due to sedation.

• In the Evening Before the Procedure:

o Do not have anything to eat or drink after midnight.

• On the Day Of the Procedure:

- Nothing to eat or drink before procedure (no candy, cough drops, gum, chewing tobacco or snuff).
 The exception is your necessary medications (heart, blood pressure, seizure, anxiety, etc) with a small sip of water.
- o Bring your completed history form and current list of medication.

• Medication Concerns:

- o If you take <u>BLOOD THINNERS</u> like <u>warfarin (Coumadin</u>), apixaban (<u>Eliquis</u>), rivaroxaban (<u>Xarelto</u>), or others, you will need to contact your prescribing doctor to ask if you can hold this medication prior to your procedure. In general, Coumadin needs to be held for 5 days (need to make arrangements to have clotting tests (PT/INR) done prior to your procedure. This can be done the day before your test), Eliquis 2 days, and Xarelto 2 days. Baby aspirin does not need to be held.
- o If you are diabetic and require <u>insulin</u> or <u>diabetic medicine</u>, your dose may need to be adjusted because of the dietary changes associated with this test. You will need to contact your doctor that orders your medications for directions.
- o If you take certain injections for diabetes or weight loss such as semaglutide (**Ozempic**, **Wegovy**), tirzepatide (**Mounjaro**, **Zepbound**), or dulaglutide (**Trulicity**), you will need to hold dose 7 days prior to your exam (you will need to contact your prescribing doctor if you can hold it).